

Literature review

## Reconditioning in patients with rheumatoid arthritis

M.-A. Mayoux Benhamou

*Service de rééducation et de réadaptation de l'appareil locomoteur et des pathologies du Rachis, hôpital Cochin (APHP), université Paris Descartes, 27, rue du Faubourg-Saint-Jacques, 75014 Paris, France*

Received 15 February 2007; accepted 20 March 2007

### Abstract

Rheumatologists traditionally have recommended to rheumatoid arthritis (RA) patients that they avoid dynamic and weight-bearing exercises because of concerns about aggravating joint inflammation and accelerating joint damage in such patients. These restrictions may lead to inadequate levels of physical activity and deconditioning.

*Objective.* – To review the literature on tolerance and benefits of conditioning training, including dynamic and weight-bearing activities in RA patients.

*Materials and methods.* – Medline and Cochrane databases were searched with the keywords RA, rehabilitation, physical therapy, exercise, reconditioning, and rest.

*Results.* – Rest therapy is more deleterious than beneficial in most patients with RA and may lead to deconditioning. Dynamic and aerobic exercises do not aggravate joint inflammation and do not accelerate joint damage in such patients. The important goal of reconditioning patients with RA is the prevention of functional decline. Conditioning programs designed to prevent widespread morbidities in healthy subjects are attainable by most RA patients, but an individualized approach to exercise is required.

*Conclusion.* – RA patients need to be persuaded about the effectiveness and safety of moderate and even high-intensity exercise.

© 2007 Elsevier Masson SAS. All rights reserved.

*Keywords:* Rheumatoid arthritis; Rehabilitation; Physical therapy; Exercise; Reconditioning; Rest

### 1. Introduction

Rheumatoid arthritis (RA) is a systemic, chronic inflammatory rheumatism that leads to joint pain, deformity and destruction of joints. Patients with RA are at risk of decreased physical capacity and functional ability. Consequently, they have increased susceptibility for comorbid conditions such as cardiovascular disease or osteoporosis [14,34,39].

Regular exercise is encouraged to prevent such widespread morbidities. For people with RA, aerobic rather than nonaerobic exercise appears to be more beneficial [24] and dynamic exercise, requiring muscle work during joint motion, superior to static or isometric exercise [7]. Nevertheless, rheumatologists traditionally have recommended that RA patients exercise restriction or that exercise programs be limited to nonweight-

bearing isometric exercises and range-of-motion exercises because of concerns about aggravating joint inflammation and accelerating joint damage in such patients [16,23,33]. These restrictions in intensity, frequency and/or duration of exercise may lead to inadequate levels of physical activity and deconditioning [32].

Since 1985 [13], a large number of controlled and uncontrolled studies have been conducted to question whether patients with RA could benefit from regular physical exercise, including dynamic and weight-bearing exercises with moderate- to high-intensity exercise programs. The results are summarized in several reviews on exercise and RA [6,12,25,31,36,38]. The consensus of these reviews is that people with RA demonstrate improved aerobic fitness, muscle strength, joint mobility, functional ability, and mood with exercise, without detrimental effects on disease. Consequently, the American College of Rheumatology (ACR) recommended regular participation in dynamic exercise programs in its recent update of treatment guidelines for RA [8]. We aimed to review the litera-

*E-mail address:* [anne.mayoux-benhamou@cch.aphp.fr](mailto:anne.mayoux-benhamou@cch.aphp.fr)  
(M.-A. Mayoux Benhamou).

ture on reconditioning in RA patients to answer specific questions related to the topic.

## 2. Methods

### 2.1. Questions

The issue of reconditioning in RA patients prompted us to formulate several questions:

- Is rest therapy deleterious or beneficial?
- Do dynamic and aerobic exercises increase joint inflammation?
- Do dynamic and aerobic exercises accelerate joint damage?
- What are the goals of reconditioning?

Are conditioning programs designed to prevent widespread morbidities, in particular cardiovascular disease, attainable by patients with RA?

### 2.2. Literature review strategy

We searched the Medline and Cochrane databases using the keywords RA, rehabilitation, physical therapy, exercise, reconditioning, and rest. Publications written in English or French selected for the review were meta-analyses, reviews (systematic or not), results of randomized controlled trials, or results of randomized uncontrolled trials. We also conducted a manual search of the references listed in each publication.

### 2.3. Patients

Only reports on adult patients with RA defined by the ACR criteria were selected.

## 3. Results

### 3.1. Rest therapy is more deleterious than beneficial in most patients

On the basis of the simplistic hypothesis that the pressure and frictional stress produced by exercise contributes to joint destruction in RA [15], bed rest and/or reduction of the exercise load have usually been recommended for patients with RA. However, in a controlled clinical trial, the benefit of bed rest was less than expected, and the positive response to rest was confined to patients with the most active disease [1]. Likewise, a substantial reduction in intensity, frequency and/or duration of exercise may lead to deconditioning [32], which is deleterious. Not exercising enough could cause a downward spiral in the condition of the patient: inactivity causes isolation, which produces anxiety and depression, which increases pain levels, leading to more inactivity. Thus, empowering RA patients through exercise could disrupt this spiral.

Rest therapy proposed for numerous rheumatic diseases is currently questionable [2], and the reinterpretation of pain by

use of vigorous exercise despite pain has been used successfully for patients with chronic pain. Such a reconditioning program could be beneficial for RA patients, although exercise-induced pain might be an indicator of an active disease process.

### 3.2. Dynamic and aerobic exercises do not aggravate joint inflammation

Numerous clinical trials of exercise concluded that moderate- to high-intensity exercises, including dynamic exercises and weight-bearing aerobic activities, do not negatively affect the disease process in people with relatively stable RA. Three studies on this topic are particularly interesting [6]. de Jong et al. [3] compared the effectiveness and safety of a 2-year intensive weight-bearing exercise program in a randomized controlled trial. Three hundred patients with stable disease were enrolled. Subjects allocated to the intensive exercise program exercised in groups twice a week for 75 min per session using the Rheumatoid Arthritis Patients in Training program (RAPIT program). Subjects allocated to the usual care group received physical therapy if thought necessary by the attending physician. In both groups, disease activity, as measured by the Disease Activity Score (DAS), decreased, with no differences between the two groups.

Häkkinen et al. [11] designed a randomized controlled trial comparing the long-term effects of a 2-year home-based strength-training program with conventional training in patients with early RA. The program consisted of dynamic strength-training exercises for the limbs and trunk to be performed twice a week for 45 min. In addition, subjects were encouraged to participate in recreational sports activities two to three times a week. The conventional training program consisted of stretching exercises performed twice a week. Compared with the baseline DAS, disease activity decreased in both groups, with even a significant difference between the two groups in favor of the strength-training group. After 5 years, both groups exhibited decreased disease activity [10]. Van den Ende et al. [37] examined the effects of intensive exercise in patients with active RA. In this randomized controlled trial, patients were allocated to an intensive exercise program consisting of supervised isokinetic and isometric strength-training and bicycling on a home trainer for 15 min, three times a week, plus conservative exercises consisting of stretching and isometric exercises. Control patients received conservative exercises only. The improvement of disease activity, as measured by the DAS, over the 24-week follow-up period was similar in both groups.

### 3.3. Dynamic and aerobic exercises do not accelerate joint damage

A few well-designed studies assessed the long-term effects of high-intensity exercises on joint damage as seen on radiography in patients with RA. None showed an increase in the rate of damage following dynamic exercise programs.

In a 5-year follow-up randomized controlled trial, Nordemar et al. [28] assessed joint damage using radiographs of symptomatic joints of the feet and based on the Larsen scoring method. Patients in the exercise group participated in group training sessions conducted by a physical therapist for 60 min once a week. The exercise regimen included various aerobic and weight-bearing activities such as cycling, dancing, or jogging. The authors reported no radiographic evidence of pronounced progression in the exercise group following a regular, supervised exercise program as compared with the control group matched in age, disease duration, and disease activity. Häkkinen et al., in the trial of the 2-year program, did not detect a significant difference in joint damage between the groups undergoing a strength-training or conventional training program, although joint damage increased slightly in both groups at 2-year follow-up [11] and at 5-year follow-up [10]. de Jong et al. [5] concluded that high-intensity weight-bearing exercises included in the RAPIT program did not increase damage of the joints of the hands and feet as seen on radiography and might even retard the progression of joint damage in the feet. However, the safety regarding the impact of high-intensity exercise on large joints with pre-existing joint damage is still under debate.

#### *3.4. The goal of reconditioning in patients with RA is the prevention of functional decline*

Dynamic exercises improve muscle function and aerobic capacity [3,7,9–11,24,28,30,35,37] and functional ability [3, 10,11,35], but exercises of at least moderate intensity, for 30 min/day for 5 days or more of the week, are necessary. This level of intensity corresponds with cardiovascular prevention programs recommended for the general population [21]. To slow age-related bone loss, impact-generating activities seem essential [17,18], or strength-training, with effects specific to the sites loaded [22]. The RAPIT program was effective in slowing the age-related loss of bone mineral density of the hip [4], although a 5-year intensive home-based muscle strength-training was not able to stop bone density loss in the Häkkinen et al. [10] study.

The psychological benefit of exercise in RA patients has been sparsely studied. Regular participation in moderate aerobic exercise may improve emotional status [9], alleviate fatigue [13,27,29] and improve quality of life [4,24].

#### *3.5. Thus, conditioning programs designed to prevent widespread morbidities in healthy subjects are attainable by most RA patients, but an individualized approach to exercise is required*

Although, low-intensity land-based or water-based exercises can be advised safely for patients with RA at any stage of the disease, a tailor-made program, partly supervised at least, could make it possible for these patients to participate in dynamic, weight-bearing and moderate- to high-intensity exercises, either individually or with group programs. Likewise, because

of the variable course of RA, the patient's ability to modify the exercise program according to current health status and responses to the exercises is very important [11]. Patients with relatively stable RA and no extensive involvement of the large joints who understand and recognize their opportunities and limitations might be able to join community-based physical activity programs, with initial or periodic supervision [6].

Finally, conditioning training, including moderate- or high-intensity and weight-bearing exercises, are recommended for most patients with RA to improve their physical health and psychological well-being and do not have a detrimental effect on disease activity or progression of joint damage as seen on radiography. However, a recent report stated that most rheumatologists and physical therapists still recommend conventional RA exercise programs with exercise restrictions [26] and that half of the patients who receive an exercise prescription are referred to physical therapy [19] with its questionable cost-effectiveness [20]. Thus, disseminating knowledge about the effectiveness and safety of moderate- and even high-intensity exercise for RA patients is needed, as is persuading health care to modify RA medical care.

#### **References**

- [1] Alexander GJM, Hortas C, Bacon PA. Bed rest, activity and the inflammation of the rheumatoid arthritis. *Br J Rheumatol* 1983;22:134–40.
- [2] Allen C, Gasziou P, Del Mar C. Bed rest: a potential harmful treatment needing more careful evaluation. *Lancet* 1999;354:1229–33.
- [3] de Jong Z, Munneke M, Zwiderman AH, Kroon HM, Jansen A, Ronda KH, et al. Is a long-term high-intensity exercise program effective and safe in patients with rheumatoid arthritis? Results of a randomized controlled trial. *Arthritis Rheum* 2003;48:2415–24.
- [4] de Jong Z, Munneke M, Lems WF, Zwiderman AH, Kroon HM, Pauwels EK, et al. Slowing of bone loss in patients with rheumatoid arthritis by long-term high-intensity exercise: results of a randomized, controlled trial. *Arthritis Rheum* 2004;50:1066–76.
- [5] de Jong Z, Munneke M, Zwiderman AH, Kroon KH, Ronda KH, Lems WF, et al. Long term high intensity exercise and damage of small joints in rheumatoid arthritis. *Ann Rheum Dis* 2004;63:1399–405.
- [6] de Jong Z, Vliet Vlieland TPM. Safety of exercise in patients with rheumatoid arthritis. *Curr Opin Rheumatol* 2005;17:177–82.
- [7] Ekdahl C, Andersson SI, Moritz U, et al. Dynamic versus static training in patients with rheumatoid arthritis. *Scand J Rheumatol* 1990;19:17–26.
- [8] Guidelines for the management of rheumatoid arthritis: 2002 update. *Arthritis Rheum* 2002;46:328–46.
- [9] Häkkinen A, Häkkinen K, Hannonen P. Effects of strength training on neuromuscular function and disease activity in patients with recent-onset inflammatory arthritis. *Scand J Rheumatol* 1994;23:237–42.
- [10] Häkkinen A, Sokka T, Kautiainen H, Kotaniemi A, Hannonen P. Sustained maintenance of exercise induced muscle strength gains and normal bone mineral density in patients with early rheumatoid arthritis: a 5 year follow up. *Ann Rheum Dis* 2004;63:910–6.
- [11] Häkkinen A, Sokka T, Kotaniemi A, Hannonen P. A randomized two-year study of the effects of dynamic strength training on muscle strength, disease activity, functional capacity, and bone mineral density in early rheumatoid arthritis. *Arthritis Rheum* 2001;44:515–22.
- [12] Häkkinen A. Effectiveness and safety of strength training in rheumatoid arthritis. *Curr Opin Rheumatol* 2004;16:132–7.
- [13] Harkcom TM, Lampman RM, Banwell BF, Castor CW. Therapeutic value of graded aerobic exercise training in rheumatoid arthritis. *Arthritis Rheum* 1985;28:32–9.

- [14] Haugeberg G, Orstavik RE, Uhlig T, Falch JA, Halse JI, Kvien TK. Bone loss in patients with rheumatoid arthritis: results from a population-based cohort of 366 patients followed up for two years. *Arthritis Rheum* 2002;46:1720–8.
- [15] Jayson MIV, Dixon AS, Yeoman P. Usual geodes [bone cysts] in rheumatoid arthritis. *Ann Rheum Dis* 1972;31:174–8.
- [16] Jivoff L. Rehabilitation and rheumatoid arthritis. *Bull Rheum Dis* 1975;26:838–41.
- [17] Kelley GA, Kelley KS, Tran ZV. Exercise and bone mineral density in men: a meta-analysis. *J Appl Physiol* 2000;88:1730–6.
- [18] Kelley GA, Kelley KS, Tran ZV. Resistance training and bone mineral density in women: a meta-analysis of controlled trials. *Am J Phys Med Rehabil* 2001;80:65–77.
- [19] Li LC, Bombardier C. Utilization of physiotherapy and occupational therapy by Ontario rheumatologists in managing rheumatoid arthritis—a survey. *Physiother Can* 2003;55:23–30.
- [20] Li LC, Iversen MD. Outcomes of patients with rheumatoid arthritis receiving rehabilitation. *Curr Opin Rheumatol* 2005;17:172–6.
- [21] Manson JE, Greenland P, LaCroix AZ, Stefanick ML, Mouton CP, Oberman A, et al. Walking compared with vigorous exercise for the prevention of cardiovascular events in women. *N Engl J Med* 2002;347:716–25.
- [22] Mayoux-Benhamou MA, Leyge JF, Roux C, Revel M. Cross sectional study on site-specific effect of weight bearing activity on proximal femur bone density. *Calcif Tissue Int* 1999;64:179–83.
- [23] Mills J, Pinals R, Ropes M, Short CL, Sutcliffe J. Value of bed rest in patients with rheumatoid arthritis. *N Engl J Med* 1971;284:453–8.
- [24] Minor MA, Hewett JE, Webel RR, Anderson SK, Kay DR. Efficacy of physical conditioning exercise in patients with rheumatoid arthritis and osteoarthritis. *Arthritis Rheum* 1989;82:1396–405.
- [25] Munneke M, de Jong Z. The role of exercise programs in the rehabilitation of patients with rheumatoid arthritis. *Int Sport Med J* 2000;1:12.
- [26] Munneke M, de Jong Z, Zwinderman AH, Runday HK, Van den Ende CH, Vliet Vlieland TP, et al. High intensity exercise or conventional exercise for patients with rheumatoid arthritis? Outcome expectations of patients, rheumatologists, and physiotherapists. *Ann Rheum Dis* 2004;63:804–8.
- [27] Neuberger GB, Press AN, Lindsley HB, Hinton R, Cagle PE, Carlson K, et al. Effects of exercise on fatigue, aerobic fitness, and disease activity measures in persons with rheumatoid arthritis. *Res Nurs Health* 1997;20:195–204.
- [28] Nordemar R, Ekblom B, Zachrisson L, Lundqvist K. Physical training in rheumatoid arthritis. A controlled long-term study, I. *Scand J Rheumatol* 1981;10:17–23.
- [29] Perlman SG, Connell KJ, Clark A, Robinson MS, Conlon P, Gecht M, et al. Dance-based aerobic exercise for rheumatoid arthritis. *Arthritis Care Res* 1990;3:29–35.
- [30] Stenström CH, Lindell B, Swanberg E, Swanberg P, Harms-Ringdahl K, Nordemar R. Intensive dynamic training in water for rheumatoid arthritis functional class II—a long-term study of effects. *Scand J Rheumatol* 1991;20:358–65.
- [31] Stenström CH, Minor MA. Evidence for the benefit of aerobic and strengthening exercise in rheumatoid arthritis. *Arthritis Rheum* 2003;49:428–34.
- [32] Stewart AL, Hays RD, Wells KB, Rogers WH, Spritzer KL, Greenfield S. Long-term functioning and well-being outcomes associated with physical activity and exercise in patients with chronic conditions in the Medical Outcomes Study. *J Clin Epidemiol* 1994;47:719–30.
- [33] Swezey R. Essentials of physical management and rehabilitation in arthritis. *Semin Arthritis Rheum* 1974;3:349–68.
- [34] Turesson C, Jarenros A, Jacobson L. Increased incidence of cardiovascular disease in patients with rheumatoid arthritis: results from a community based study. *Ann Rheum Dis* 2004;63:952–5.
- [35] van den Ende CH, Hazes JM, le Cessie S, Mulder WJ, Belfor DG, Breedveld FC, et al. Comparison of high and low intensity training in well controlled rheumatoid arthritis: results of a randomized clinical trial. *Ann Rheum Dis* 1996;55:798–805.
- [36] van den Ende CH, Vliet Vlieland TP, Munneke M, Hazes JM. Dynamic exercise therapy in rheumatoid arthritis: a systematic review. *Br J Rheumatol* 1998;37:677–87.
- [37] van den Ende CH, Breedveld FC, le Cessie S, Dijkmans BA, de Mug AW, Hazes JM. Effect of intensive exercise on patients with active rheumatoid arthritis: a randomized clinical trial. *Ann Rheum Dis* 2000;59:615–21.
- [38] Westby MD. A health professional's guide to exercise prescription for people with arthritis: a review of aerobic fitness activities. *Arthritis Care Res* 2001;45:501–10.
- [39] Wong JB, Ramey DR, Singh G. Long-term morbidity, mortality, and economics of rheumatoid arthritis. *Arthritis Rheum* 2001;44:2746–9.